arXiv:1411.7205v1 [math.RA] 26 Nov 2014

Relative Hom-Hopf modules and total integrals

Shuangjian Guo[1], Xiaohui Zhang[2] and Shengxiang Wang[3*]

1. School of Mathematics and Statistics, Guizhou University of Finance and Economics, Guizhou 550025, China

2. Department of Mathematics, Southeast University, Nanjing 210096, China

3. School of Mathematics and Finance, Chuzhou University, Chuzhou 239000, China

Abstract Let (H,α) be a monoidal Hom-Hopf algebra and (A,β) a right (H,α)-Hom-comodule algebra. We first investigate the criterion for the existence of a total integral of (A,β) in the setting of monoidal Hom-Hopf algebras. Also we prove that there exists a total integral $\phi:(H,\alpha)\to(A,\beta)$ if and only if any representation of the pair (H,A) is injective in a functorial way, as a corepresentation of (H,α), which generalizes Doi's result. Finally, we define a total quantum integral $\gamma: H\to Hom(H,A)$ and prove the following affineness criterion: if there exists a total quantum integral γ and the canonical map $\psi: A\otimes_B A\to A\otimes H,\ \ a\otimes_B b\mapsto \beta^{-1}(a)b_{[0]}\otimes\alpha(b_{[1]})$ is surjective, then the induction functor $A\otimes_B -:\widetilde{\mathscr{H}}(\mathscr{M}_k)_B\to\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ is an equivalence of categories.

Keywords Monoidal Hom-Hopf algebra; total integral; relative Hom-Hopf module.

MR(2010)Subject Classification 16T05

1 Introduction

Hom-algebras and Hom-coalgebras were introduced by Makhlouf and Silvestrov in [14] as generalizations of ordinary algebras and coalgebras in the following sense: the associativity of the multiplication is replaced by the Hom-associativity and similar for Hom-coassociativity. They also defined the structures of Hom-bialgebras and Hom-Hopf algebras, and described some of their properties extending properties of ordinary bialgebras and Hopf algebras in [15] and [16]. Recently, many more properties and structures of Hom-Hopf algebras have been developed, see [3], [9], [12], [22] and references cited therein.

Caenepeel and Goyvaerts [1] studied Hom-bialgebras and Hom-Hopf algebras from a categorical view point, and called them monoidal Hom-bialgebras and monoidal Hom-Hopf algebras respectively, which are slightly different from the above Hom-bialgebras and Hom-Hopf algebras. In [2], Chen, Wang and Zhang introduced integrals of monoidal Hom-Hopf algebras and investigated the existence and uniqueness of integrals for finite-dimensional monoidal Hom-Hopf algebras, they also applied the integrals to the Maschke

*Correspondence: S X Wang: wangsx-math@163.com.

type theorem for monoidal Hom-Hopf algebras controlling the semisimplicity and separability of monoidal Hom-Hopf algebras.

In [12], the first author and Chen introduced the notion of relative Hom-Hopf modules and proved that the functor F from the category of relative Hom-Hopf modules to the category of right (A,β)-Hom-modules has a right adjoint. We also introduced the notion of total integrals for Hom-comodule algebras which has effective applications in investigating properties of relative Hom-Hopf modules, and proved a Maschke type theorem for the category of relative Hom-Hopf modules as an application our theory.

Total integral is a vital notion in representation theories. Doi [7] introduced the notion of total integral for any H-comodule algebra A over a Hopf algebra H, which has strong ties both to $\mathcal{M}^H$ (i. e., the corepresentation of H) and to the representation of the pair (H,A) (i. e., the category of relative Hopf modules $\mathcal{M}_A^H$), and presented the famous necessary and sufficient criterion for the existence of a total integral. Later, Menini and Militaru [17] interpreted the criterion for the existence of a total integral with the help of forgetful functors $F:\mathcal{M}_A^H\to\mathcal{M}_A$ and $G:\mathcal{M}_A^H\to\mathcal{M}^H$. Furthermore, the affineness criterion for affine algebraic group schemes was studied by Cline et al. [5] and Oberst [18]. Inspired by the ideas from Doi, Menini and Militaru, we want to investigate the criterion for the existence of a total integral in the setting of monoidal Hom-Hopf algebras.

The paper is organized as follows. In Section 2, we recall some definitions and properties relative to separable functors and monodial Hom-Hopf algebras which are needed later. In Section 3, we present two pairs of important adjoint functors (see Proposition 3.1). In Section 4, we show a necessary and sufficient criterion for the existence of an integral on a monodial Hom-Hopf algebra (see Theorem 4.3). In Section 5, we give the affineness criterion for relative Hom-Hopf modules (see Theorem 5.7).

2 Preliminaries

In this section we recall some basic definitions and results related to our paper. Throughout the paper, all algebraic systems are supposed to be over a commutative ring k. The reader is referred to [1] and [12] as general references about Hom-structures.

Let $\mathcal{C}$ be a category, there is a new category $\widetilde{\mathscr{H}}(\mathcal{C})$ as follows: objects are couples (M,μ), with $M\in\mathcal{C}$ and $\mu\in Aut_{\mathcal{C}}(M)$. A morphism $f:(M,\mu)\to(N,\nu)$ is a morphism $f:M\to N$ in $\mathcal{C}$ such that $\nu\circ f=f\circ\mu$.

Let $\mathscr{M}_k$ denotes the category of k-modules. $\mathscr{H}(\mathscr{M}_k)$ will be called the Hom-category associated to $\mathscr{M}_k$. If $(M,\mu)\in\mathscr{M}_k$, then $\mu:M\to M$ is obviously a morphism in $\mathscr{H}(\mathscr{M}_k)$. It is easy to show that $\widetilde{\mathscr{H}}(\mathscr{M}_k)=(\mathscr{H}(\mathscr{M}_k),\otimes,(I,I),\widetilde{a},\widetilde{l},\widetilde{r})$ is a monoidal category by Proposition 1.1 in [1]: the tensor product of (M,μ) and (N,ν) in $\widetilde{\mathscr{H}}(\mathscr{M}_k)$ is given by the formula $(M,\mu)\otimes(N,\nu)=(M\otimes N,\mu\otimes\nu)$.

Assume that $(M,\mu),(N,\nu),(P,\pi)\in\widetilde{\mathscr{H}}(\mathscr{M}_k)$. The associativity and unit constraints are given by the formulas

$$\widetilde{a}_{M,N,P}((m\otimes n)\otimes p)=\mu(m)\otimes(n\otimes\pi^{-1}(p)),$$
$$\widetilde{l}_M(x\otimes m)=\widetilde{r}_M(m\otimes x)=x\mu(m).$$

An algebra in $\widetilde{\mathscr{H}}(\mathscr{M}_k)$ will be called a monoidal Hom-algebra.

Definition 2.1. A monoidal Hom-algebra is an object $(A,\alpha)\in\widetilde{\mathscr{H}}(\mathscr{M}_k)$ together with a k-linear map $m_A: A\otimes A\to A$ and an element $1_A\in A$ such that

$$\begin{aligned}&\alpha(ab)=\alpha(a)\alpha(b);\ \alpha(1_A)=1_A,\\&\alpha(a)(bc)=(ab)\alpha(c);\ a1_A=1_Aa=\alpha(a),\end{aligned}$$

for all $a,b,c\in A$. Here we use the notation $m_A(a\otimes b)=ab$.

Definition 2.2. A monoidal Hom-coalgebra is an object $(C,\gamma)\in\widetilde{\mathscr{H}}(\mathscr{M}_k)$ together with k-linear maps $\Delta: C\to C\otimes C,\quad \Delta(c)=c_{(1)}\otimes c_{(2)}$ (summation implicitly understood) and $\gamma: C\to C$ such that

$$\Delta(\gamma(c))=\gamma(c_{(1)})\otimes\gamma(c_{(2)});\quad \varepsilon(\gamma(c))=\varepsilon(c),$$

and

$$\begin{aligned}&\gamma^{-1}(c_{(1)})\otimes c_{(2)(1)}\otimes c_{(2)(2)}=c_{(1)(1)}\otimes c_{(1)(2)}\otimes\gamma(c_{(2)}),\\&\varepsilon(c_{(1)})c_{(2)}=\varepsilon(c_{(2)})c_{(1)}=\gamma^{-1}(c),\end{aligned}$$

for all $c\in C$.

Definition 2.3. A monoidal Hom-bialgebra $H=(H,\alpha,m,\eta,\Delta,\varepsilon)$ is a bialgebra in the category $\widetilde{\mathscr{H}}(\mathscr{M}_k)$. This means that (H,α,m,η) is a monoidal Hom-algebra, (H,Δ,α) is a monoidal Hom-coalgebra and that Δ and ε are morphisms of monoidal Hom-algebras, that is,

$$\begin{aligned}&\Delta(ab)=a_{(1)}b_{(1)}\otimes a_{(2)}b_{(2)};\Delta(1_H)=1_H\otimes 1_H,\\&\varepsilon(ab)=\varepsilon(a)\varepsilon(b),\ \varepsilon(1_H)=1_H.\end{aligned}$$

Definition 2.4. A monoidal Hom-Hopf algebra is a monoidal Hom-bialgebra (H,α) together with a linear map $S: H\to H$ in $\widetilde{\mathscr{H}}(\mathscr{M}_k)$ such that

$$S*I=I*S=\eta\varepsilon,\ S\alpha=\alpha S.$$

Definition 2.5. Let (A,α) be a monoidal Hom-algebra. A right (A,α)-Hom-module is an object $(M,\mu)\in\widetilde{\mathscr{H}}(\mathscr{M}_k)$ consists of a k-module and a linear map $\mu: M\to M$ together with a morphism $\psi: M\otimes A\to M, \psi(m\cdot a)=m\cdot a$ in $\widetilde{\mathscr{H}}(\mathscr{M}_k)$ such that

$$(m\cdot a)\cdot\alpha(b)=\mu(m)\cdot(ab);\qquad m\cdot 1_A=\mu(m),$$

for all $a\in A$ and $m\in M$. The fact that $\psi\in\widetilde{\mathscr{H}}(\mathscr{M}_k)$ means

$$\mu(m\cdot a)=\mu(m)\cdot\alpha(a).$$

A morphism $f: (M,\mu)\to(N,\nu)$ in $\widetilde{\mathscr{H}}(\mathscr{M}_k)$ is called right A-linear if it preserves the A-action, that is, $f(m\cdot a)=f(m)\cdot a$. $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A$ will denote the category of right (A,α)-Hom-modules and A-linear morphisms.

Definition 2.6. Let (C,γ) be a monoidal Hom-coalgebra. A right (C,γ)-Hom-comodule is an object $(M,\mu)\in\widetilde{\mathscr{H}}(\mathscr{M}_k)$ together with a k-linear map $\rho_M: M\to M\otimes C$ $(\rho_M(m)=m_{[0]}\otimes m_{[1]})$ in $\widetilde{\mathscr{H}}(\mathscr{M}_k)$ such that

$$m_{[0][0]}\otimes(m_{[0][1]}\otimes\gamma^{-1}(m_{[1]}))=\mu^{-1}(m_{[0]})\otimes\Delta_C(m_{[1]});\ m_{[0]}\varepsilon(m_{[1]})=\mu^{-1}(m),$$

for all $m\in M$. The fact that $\rho_M\in\widetilde{\mathscr{H}}(\mathscr{M}_k)$ means

$$\rho_M(\mu(m))=\mu(m_{[0]})\otimes\gamma(m_{[1]}).$$

Morphisms of right (C,γ)-Hom-comodule are defined in the obvious way. The category of right (C,γ)-Hom-comodules will be denoted by $\widetilde{\mathscr{H}}(\mathscr{M}_k)^C$.

Definition 2.7. Let (H,α) be a monoidal Hom-Hopf algebra. A monoidal Hom-algebra (A,β) is called a right (H,α)-Hom-comodule algebra if (A,β) is a right (H,α) Hom-comodule with a coaction $\rho_A: A\to A\otimes H, \rho_A(a)=a_{[0]}\otimes a_{[1]}$ such that the following conditions hold:

$$\rho_A(ab)=a_{[0]}b_{[0]}\otimes a_{[1]}b_{[1]}, \rho_A(1_A)=1_A\otimes 1_H,$$

for all $a,b\in A$.

3 Relative Hom-Hopf Modules: functors and structures

Let (H,α) be a monoidal Hom-Hopf algebra and (A,β) a right (H,α)-Hom-comodule algebra. Recall from [12], a relative Hom-Hopf module (M,μ) is a right (A,β)-Hom-module which is also a right (H,α)-Hom-comodule with the coaction structure $\rho_M: M\to M\otimes H$ defined by $\rho_M(m)=m_{[0]}\otimes m_{[1]}$ such that the following compatible condition holds: for all $m\in M$ and $a\in A$,

$$\rho_M(ma)=m_{[0]}\cdot a_{[0]}\otimes m_{[1]}a_{[1]}.$$

A morphism between two right relative Hom-Hopf modules is a k-linear map which is a morphism in the categories $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A$ and $\widetilde{\mathscr{H}}(\mathscr{M}_k)^H$ at the same time. $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ will denote the category of right relative Hom-Hopf modules and morphisms between them.

By [12], the forgetful functor $F: \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H\to\widetilde{\mathscr{H}}(\mathscr{M}_k)_A$ has a right adjoint $G: \widetilde{\mathscr{H}}(\mathscr{M}_k)_A\to\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. G defined by

$$G(M)=M\otimes H$$

with structure maps

$$(m\otimes h)\cdot a=m\cdot a_{[0]}\otimes ha_{[1]},$$
$$\rho_{G(M)}(m\otimes h)=(\mu^{-1}(m)\otimes h_{(1)})\otimes\alpha(h_{(2)}),$$

for all $a\in A, m\in M$ and $h\in H$.

Let us describe the unit η and the counit δ of the adjunction. The unit is described by the coaction: for $M\in\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$, we define $\eta_M: M\to M\otimes H$ as follows:

$$\eta_M(m)=m_{[0]}\otimes m_{[1]},\ m\in M.$$

We can check that $\eta_M \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. For any $N \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A$, define

$$\delta_N : N \otimes H \to N,\ \delta_N(n \otimes h) = \varepsilon(h)n,\ n \in N, h \in H.$$

It is easy to check that $\eta_M \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. One may verify that η and δ defined above are natural transformations and satisfy

$$G(\delta_N) \circ \eta_{G(N)} = I_{G(N)},\ \delta_{F(M)} \circ F(\eta_M) = I_{F(M)},$$

for all $M \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ and $N \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A$.

Since (A, β) is a right (A, β)-Hom-module, $G(A)$ is a relative Hom-Hopf module via

$$(a \otimes h) \cdot b = ab_{[0]} \otimes hb_{[1]}; \tag{3.1}$$
$$\rho_{A\otimes H}(a \otimes h) = (\beta^{-1}(a) \otimes h_{(1)}) \otimes \alpha(h_{(2)}), \tag{3.2}$$

for all $a, b \in A$ and $h \in H$. Notice that $GFG(A) = A \otimes H \otimes H$ is an object in $\widetilde{\mathscr{H}}(\mathscr{M}_k)^H$, and we also have

$$\begin{cases} (a \otimes h \otimes g) \cdot b = ab_{[0][0]} \otimes hb_{[0][1]} \otimes gb_{[1]}; \\ \rho_{A\otimes H\otimes H}(a \otimes h \otimes g) = (\beta^{-1}(a) \otimes \alpha^{-1}(h) \otimes g_{(1)}) \otimes \alpha(g_{(2)}), \end{cases}$$

for any $a, b \in A$ and $h, g \in H$.

Similar to [12], one can check that $\widetilde{F} : \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H \to \widetilde{\mathscr{H}}(\mathscr{M}_k)^H$ which forgets the (A, β)-action has a left adjoint given by

$$\widetilde{G} : \widetilde{\mathscr{H}}(\mathscr{M}_k)^H \to \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H, \widetilde{G}(N) = A \otimes N,\ N \in \widetilde{\mathscr{H}}(\mathscr{M}_k)^H.$$

The relative Hom-Hopf module structure on $\widetilde{G}(N)$ is given by

$$\begin{cases} (a \otimes n) \cdot b = a\beta(b) \otimes \nu^{-1}(n), \\ \rho_{A\otimes N}(a \otimes n) = (a_{[0]} \otimes n_{[0]}\otimes) \otimes a_{[1]}n_{[1]}, \end{cases}$$

for all $a \in A$ and $n \in N$. Since (H, α) is a right (H, α)-Hom-comodule, then $\widetilde{G}(H) = A\otimes H$ can be viewed as a relative Hom-Hopf module via

$$(a \otimes h) \cdot b = a\beta(b) \otimes \alpha^{-1}(h), \tag{3.3}$$
$$\rho_{A\otimes H}(a \otimes h) = (a_{[0]} \otimes h_{(1)}\otimes) \otimes a_{[1]}h_{(2)}, \tag{3.4}$$

for all $a \in A$ and $h \in H$.

From Eqs. (3.1) and (3.2) or Eqs. (3.3) and (3.4), we have two types of relative Hom-Hopf modules structures on $A \otimes H$. The following proposition reveals their relationship.

Proposition 3.1. With the two types of relative Hom-Hopf modules on $A \otimes H$. Then

$$G(A) \cong \widetilde{G}(H)$$

in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$.

Proof. For any $a \in A$ and $h \in H$, we construct two maps u and v as follows:

$$u : A \otimes H \to A \otimes H, u(a \otimes h) = \beta(a_{[0]}) \otimes h\alpha^{-1}(a_{[1]}),$$
$$v : A \otimes H \to A \otimes H, u(a \otimes h) = \beta(a_{[0]}) \otimes h\alpha^{-1}(S(a_{[1]})).$$

We claim that $u \circ v = id_{A\otimes H}$. In fact,

$$\begin{aligned} u \circ v(a \otimes h) &= u(\beta(a_{[0]}) \otimes h\alpha^{-1}(S(a_{[1]}))) \\ &= \beta^2(a_{[0][0]}) \otimes (h\alpha^{-1}(S(a_{[1]})))\alpha^{-1}(a_{[0][1]}) \\ &= \beta(a_{[0]}) \otimes (hS(a_{[1](2)}))\alpha^{-1}(a_{[1](1)}) \\ &= \beta(a_{[0]}) \otimes \alpha^{-1}(h)(S(a_{[1](2)})a_{[1](1)}) \\ &= \beta(a_{[0]}) \otimes \alpha^{-1}(h)\varepsilon(a_{[1]}) \\ &= a \otimes h. \end{aligned}$$

Similarly, one may verify that $v \circ u = id_{A\otimes H}$. The proof that u is a morphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ is straightforward. □

4 Natural transformations versus integrals

Definition 4.1.[12] Let (H, α) be a monoidal Hom Hopf algebra and (A, β) a right (H, α)-Hom-comodule algebra. The map $\phi : (H, \alpha) \to (A, \beta)$ is called a total integral if the following conditions are satisfied:

$$\rho_A\phi = (\phi \otimes id_H)\Delta_H, \quad \phi\alpha = \beta\phi, \qquad \phi(1_H) = 1_A.$$

Proposition 4.2. Let (A, β) be a right (H, α)-Hom-comodule algebra, $(M, \mu), (N, \nu) \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ and $u : (N, \nu) \to (M, \mu)$ a k-linear map. Suppose that there exists a total integral $\phi : (H, \alpha) \to (A, \beta)$. Then

(1) The map

$$\widetilde{u} : (N, \nu) \to (M, \mu),\ \widetilde{u}(n) = u(\nu(n_{[0]}))_{[0]} \cdot \phi(S(u(n_{[0]})_{[1]})\alpha^{-1}(n_{[1]})),\ n \in N$$

is right (H, α)-colinear.

(2) If $f : (M, \mu) \to (N, \nu)$ is a morphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ of which is a k-split injection (respectively, a k-split surjection), then f has an (H, α)-colinear retraction (respectively, a section).

Proof. For any $n \in N$, we have

$$\begin{aligned}
\rho_M \circ \widetilde{u}(n) &= u(\nu(n_{[0]}))_{[0][0]} \cdot \phi(S(u(n_{[0]})_{[1]})\alpha^{-1}(n_{[1]}))_{[0]} \\
&\qquad \otimes u(\nu(n_{[0]}))_{[0][1]}\phi(S(u(n_{[0]})_{[1]})\alpha^{-1}(n_{[1]}))_{[1]} \\
&= u(n_{[0]})_{[0]} \cdot \phi(S(u(\nu(n_{[0]}))_{[1](2)})\alpha^{-1}(n_{[1]}))_{[0]} \\
&\qquad \otimes u(\nu(n_{[0]}))_{[1](1)}\phi(S(u(\nu(n_{[0]}))_{[1](2)})\alpha^{-1}(n_{[1]}))_{[1]} \\
&= u(n_{[0]})_{[0]} \cdot \phi(S(u(\nu(n_{[0]}))_{[1](2)(2)})\alpha^{-1}(n_{[1](1)})) \\
&\qquad \otimes u(\nu(n_{[0]}))_{[1](1)}S(u(\nu(n_{[0]}))_{[1](2)(1)})\alpha^{-1}(n_{[1](2)}) \\
&= u(n_{[0]})_{[0]} \cdot \phi(S(u(n_{[0]})_{[1](2)})\alpha^{-1}(n_{[1](1)})) \\
&\qquad \otimes u(\nu^2(n_{[0]}))_{[1](1)(1)}S(u(\nu(n_{[0]}))_{[1](1)(2)})\alpha^{-1}(n_{[1](2)}) \\
&= u(n_{[0]})_{[0]} \cdot \phi(S(u(n_{[0]})_{[1](2)})\alpha^{-1}(n_{[1](1)})) \\
&\qquad \otimes [u(\nu(n_{[0]}))_{[1](1)(1)}S(u(\nu(n_{[0]}))_{[1](1)(2)})]n_{[1](2)} \\
&= u(n_{[0]})_{[0]} \cdot \phi(S(u(\nu^{-1}(n_{[0]}))_{[1]})\alpha^{-1}(n_{[1](1)})) \otimes \alpha(n_{[1](2)}) \\
&= u(\nu(n_{[0][0]}))_{[0]} \cdot \phi(S(u(n_{[0][0]})_{[1]})\alpha^{-1}(n_{[1](1)})) \otimes n_{[1](2)} \\
&= (\widetilde{u} \otimes id_H) \circ \rho_N(n).
\end{aligned}$$

Hence $\widetilde{u}$ is right (H, α)-colinear.

Let $u : (N, \nu) \to (M, \mu)$ be k-linear, then $\widetilde{u} : (N, \nu) \to (M, \mu)$ is right (H, α)-colinear. Assume that u is a retraction of f, for any $m \in M$, we have

$$\begin{aligned}
(\widetilde{u} \circ f)(m) &= u(\nu(f(m)_{[0]}))_{[0]} \cdot \phi(S(u(f(m)_{[0]})_{[1]})\alpha^{-1}(f(m)_{[1]})) \\
&= u(\nu(f(m_{[0])}))_{[0]} \cdot \phi(S(u(f(m_{[0]}))_{[1]})\alpha^{-1}(m_{[1]})) \\
&= \nu(m_{[0][0]}) \cdot \phi(S(m_{[0][1]})\alpha^{-1}(m_{[1]})) \\
&= m_{[0]} \cdot \phi(S(m_{[1](1)})m_{[1](2)}) \\
&= m_{[0]}\varepsilon(m_{[1]}) \cdot 1_A = m.
\end{aligned}$$

So $\widetilde{u} : (N, \nu) \to (M, \mu)$ is a right (H, α)-colinear retraction of f. On the other hand, if u is a section of f, for any $n \in N$, we have

$$\begin{aligned}
(f \circ \widetilde{u})(n) &= f(u(\nu(n_{[0]}))_{[0]}) \cdot \phi(S(u(n_{[0]})_{[1]})\alpha^{-1}(n_{[1]})) \\
&= f(u(\nu(n_{[0]}))_{[0]}) \cdot \phi(S(u(n_{[0]})_{[1]})\alpha^{-1}(n_{[1]})) \\
&= \nu(n_{[0][0]}) \cdot \phi(S(n_{[0][1]})\alpha^{-1}(n_{[1]})) \\
&= n_{[0]} \cdot \phi(S(n_{[1](1)})n_{[1](2)}) \\
&= n_{[0]}\varepsilon(n_{[1]}) \cdot 1_A = n.
\end{aligned}$$

Thus $\widetilde{u} : (N, \nu) \to (M, \mu)$ is a right H-colinear section of f. □

Theorem 4.3. Let (A, β) be a right (H, α)-Hom-comodule algebra. Then the following statements are equivalent:

(1) There exists a total integral $\phi : (H, \alpha) \to (A, \beta)$;

(2) There exists a natural transformation $\rho : \widetilde{F} \circ 1_{\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H} \to \widetilde{F} \circ G \circ F$ splits;

(3) $\rho_A : A \to A \otimes H$ splits in $\widetilde{\mathscr{H}}(\mathscr{M}_k)^H$.

Consequently, if one of the equivalent conditions holds, then any relative Hopf module is injective as a right (H,α)-comodule.

Proof. (1)⇒(2) Assume that $\phi : (H,\alpha) \to (A,\beta)$ is a total integral. We have to construct a natural transformation λ that splits ρ. Let $(M,\mu) \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ and $\varphi : M \otimes H \to M$ be the k-linear retraction of $\rho_M : M \to M \otimes H,\ \varphi(m \otimes h) = \varepsilon(h)m$. Define

$$\begin{aligned}&\lambda_M : \widetilde{F} \circ G \circ F(M) \to \widetilde{F}(M),\\ &\lambda_M(m \otimes h) = \mu(m_{[0]}) \cdot \phi(S(m_{[1]})\alpha^{-1}(h)).\end{aligned}$$

It follows from Proposition 4.2 that the map λ_M is a right (H,α)-colinear retraction of ρ_M.

It remains to prove that $\lambda_M \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ is a natural transformation. Let $f : (M,\mu) \to (N,\nu)$ be a morphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$, we have to prove that

$$f \circ \lambda_M = \lambda_N \circ (f \otimes id_H).$$

Since f is right (H,α)-colinear, we have

$$\begin{aligned}\lambda_N \circ (f \otimes id_H)(m \otimes h) &= \lambda_N(f(m) \otimes h)\\ &= \nu(f(m)_{[0]}) \cdot \phi(S(f(m)_{[1]})\alpha^{-1}(h))\\ &= \nu(f(m_{[0]})) \cdot \phi(S(m_{[1]})\alpha^{-1}(h))\\ &= f \circ \lambda_M(m \otimes h).\end{aligned}$$

So λ_M is a natural transformation that splits ρ_M.

(2)⇒(3) Assume that for any $(M,\mu) \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$, $\chi_M : \widetilde{F}(M) \to \widetilde{F} \circ G \circ F(M)$ splits in the category $\widetilde{\mathscr{H}}(\mathscr{M}_k)^H$ of right H-comodules and the character of the splitting is functorial. In particular, we consider the relative Hopf module (A,β), then we have

$$\rho_A : A \to A \otimes H, \qquad \rho_A(a) = a_{[0]} \otimes a_{[1]}$$

splits in $\widetilde{\mathscr{H}}(\mathscr{M}_k)^H$ and let

$$\lambda_A : A \otimes H \to A$$

be a right (H,α)-colinear retraction of it.

(3)⇒(1) Assume that the map $\rho_A : A \to A \otimes H, \rho_A(a) = a_{[0]} \otimes a_{[1]}$ splits in $\widetilde{\mathscr{H}}(\mathscr{M}_k)^H$. Let $\lambda_A : A \otimes H \to A$ be the map which splits ρ_A. We define

$$\phi : (H,\alpha) \to (A,\beta), \qquad \phi(h) = i_A^{-1}(\lambda_A(1_A \otimes h)),$$

where $i_A : A \to A$ is the natural isomorphism. Since λ_A is right (H,α)-colinear, we have

$$\lambda_A(1_A \otimes h_{(1)}) \otimes h_{(2)} = \lambda_A(1_A \otimes h)_{[0]} \otimes \lambda_A(1_A \otimes h)_{[1]}.$$

Let $\lambda_A(1_A \otimes h_{(1)}) = b$ and $\lambda_A(1_A \otimes h) = q$, we get $b \otimes h_{(2)} = q_{[0]} \otimes q_{[1]}$. Therefore, ϕ is (H,α)-colinear. It is sufficient to check that $\rho_A\phi = (\phi \otimes id_H)\Delta_H$. For any $h \in H$, we have

$$\rho_A \circ \phi(h) = \rho_A(i_A^{-1}(\lambda_A(1_A \otimes h))) = q_{[0]} \otimes q_{[1]}.$$

On the other hand,

$$\begin{aligned}&(\phi\otimes id_H)\Delta_H(h)=\phi(h_{(1)})\otimes h_{(2)}\\ =\ & i_A^{-1}(\lambda_A(1_A\otimes h_{(1)}))\otimes h_{(2)}=b\otimes h_{(2)}.\end{aligned}$$

Notice that $\phi(1_H)=i_A^{-1}(\lambda_A(1_A\otimes 1_H))=i_A^{-1}(\lambda_A(\rho_A(1_A)))=i_A^{-1}(1_A)=1_A$. It is easy to check that $\phi\alpha=\beta\phi$. So ϕ is a total integral. And this completes the proof. □

From Theorem 4.3, we have the following corollary.

Corollary 4.4. Let $F:\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H\to\widetilde{\mathscr{H}}(\mathscr{M}_k)_A$ and $\widetilde{F}:\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H\to\widetilde{\mathscr{H}}(\mathscr{M}_k)^H$ be two forgetful functors. Then F is $\widetilde{F}$-separable if and only if there exists a total integral $\phi:(H,\alpha)\to(A,\beta)$.

Definition 4.5. Let (H,α) be a monoidal Hom-Hopf algebra with a bijective antipode and (A,β) a right (H,α)-Hom-comodule algebra. A k-linear map $\gamma:H\to Hom(H,A)$ satisfying $\gamma(\alpha(g))(\alpha(h))=\beta\circ\gamma(g)(h)$ is called a quantum integral if

$$\gamma(\alpha^{-1}(g))(h_{(1)})\otimes\alpha(h_{(2)})=[\gamma(\alpha(g_{(2)}))(h)]_{[0]}\otimes g_{(1)}[\gamma(g_{(2)}(\alpha^{-1}(h))]_{[1]},\quad(4.\ 1)$$

for all $g,h\in H$. A quantum integral $\gamma:H\to Hom(H,A)$ is called total if

$$\gamma(h_{(1)})(h_{(2)})=\varepsilon(h)1_A,\qquad(4.\ 2)$$

for all $h\in H$.

Remark 4.6. (1) Let $\gamma:H\to Hom(H,A)$ be a quantum integral, then

$$\phi:(H,\alpha)\to(A,\beta),\ \ \phi(h)=\gamma(h)(1_H)$$

satisfies the condition:

$$\phi(h_{(1)})\otimes h_{(2)}=\phi(h)_{[0]}\otimes\phi(h)_{[1]},$$

for all $h\in H$, that is, $\phi:(H,\alpha)\to(A,\beta)$ is right (H,α)-colinear.

(2) As opposed to the case $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$, a right (H,α)-colinear map $\phi:(H,\alpha)\to(A,\beta)$ is not sufficient to construct a quantum integral. If $\phi:(H,\alpha)\to(A,\beta)$ is a k-linear map satisfying a more powerful condition:

$$g\phi(h)_{[1]}\otimes\phi(h)_{[0]}=\phi(h)_{[1]}g\otimes\phi(h)_{[0]},\ \ h,g\in H.$$

Then $\gamma:H\to Hom(H,A),\ \gamma(g)(h)=\phi(hS^{-1}(g))$ is a quantum integral.

Example 4.7. (1) Recall from that let $H=M^n(k)$ be the $n\times n$ matrix monoidal Hom-Hopf algebra with the structures given by

$$\begin{aligned}c_{iu}c_{uj}&=\delta_{ij}\alpha(c_{ij}),\qquad &\sum c_{ii}&=1_H,\\ \Delta(c_{ij})&=\alpha^{-1}(c_{iu})\otimes\alpha^{-1}(c_{uj}),\qquad &\varepsilon(c_{ij})&=\delta_{ij}.\end{aligned}$$

Let $(M^n(k),A)$ be a relative Hom-Hopf datum and $B=A^{coH}=\{a\in A|\rho(a)=\beta^{-1}(a)\otimes 1_H\}$ the subalgebra of coinvariants of A. Let $\mu=(\mu_{ij})\in M^n(B)$ be an arbitrary $n\times n$-matrix over B. Then the map

$$\gamma:H\to Hom(H,A),\ \gamma(c_{ij})(c_{rs})=\delta_{is}\beta(\mu_{rj})$$

is an integral of $(M^n(k), A)$. Indeed, for any $h = c_{ij}$ and $g = c_{kl}$, we have

$$\gamma(\alpha^{-1}(g))(h_{(1)}) \otimes \alpha(h_{(2)}) = \gamma(\alpha^{-1}(c_{kl}))(\alpha^{-1}(c_{iu})) \otimes c_{uj} = \delta_{ku}\mu_{il} \otimes c_{uj} = \mu_{il} \otimes c_{kj}$$

and

$$\begin{aligned} & [\gamma(\alpha(g_{(2)}))(h)]_{[0]} \otimes g_{(1)}[\gamma(g_{(2)}(\alpha^{-1}(h))]_{[1]} \\ = \ & [\gamma(c_{ul})(c_{ij})]_{[0]} \otimes \alpha^{-1}(c_{ku})[\gamma(\alpha^{-1}(c_{ul}))(\alpha^{-1}(c_{ij}))]_{[1]} \\ = \ & \delta_{uj}\mu_{il} \otimes c_{ku} = \mu_{il} \otimes c_{kj}. \end{aligned}$$

So γ is an integral. On the other hand, we have

$$\gamma(h_{(1)})(h_{(2)}) = \gamma(c_{iu})(c_{uj}) = \delta_{ij}\beta(\mu_{uu}).$$

Hence γ is a total integral if and only $\beta(\mu_{uu}) = 1_A$.

(2) Another class of examples of total integrals arises from the graded case. Recall from [2] that let G be a finite group and $H = kG$ be the monoidal Hom-Hopf algebra with structures:

$$\begin{aligned} & xy = \alpha(xy), 1x = \alpha(x), \varepsilon(x) = 1 \\ & \Delta(x) = \alpha^{-1}(x) \otimes \alpha^{-1}(x), \ x, y \in G. \end{aligned}$$

Let (kG, A) be a relative Hom-Hopf datum and $\mu = (\mu_{xy})_{x,y\in G}$ a family of elements of $B = A^{coH} = \{a \in A | \rho(a) = \beta^{-1}(a) \otimes 1_H\}$. Then the map

$$\gamma : H \to Hom(H, A), \ \gamma(x)(y) = \delta_{xy}\beta(\mu_{xy})$$

is an integral of (kG, A). Indeed, for any $h = x$ and $g = y$, we have

$$\begin{aligned} & [\gamma(y)(x)]_{[0]} \otimes \alpha^{-1}(y)[\gamma(\alpha^{-1}(y)(\alpha^{-1}(x))]_{[1]} \\ = \ & \delta_{xy}\mu_{xy} \otimes y = \delta_{xy}\mu_{xy}) \otimes x = \gamma(\alpha^{-1}(y))(\alpha^{-1}(x)) \otimes x. \end{aligned}$$

Furthermore, γ is a total integral iff $\mu_{xx} = 1_A$ for all $x \in G$.

Next we will give some applications of the existence of total quantum integrals.

Theorem 4.8. Let (A, β) be a right (H, α)-Hom-comodule algebra and (M, μ) an object in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. Assume that there exists a total quantum integral $\gamma : H \to Hom(H, A)$, then the map

$$\begin{aligned} & f : A \otimes H \otimes M \to M, \\ & f(a \otimes h \otimes m) = \mu(m_{[0]})\gamma(\alpha^{-1}(m_{[1]}))(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))\beta(a_{[0]}), \end{aligned}$$

for all $h \in H, a \in A$ and $m \in M$, is a k-split epimorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. In particular, $A \otimes H$ is a generator in the category $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$.

Proof. We first show that $A \otimes H \otimes M$ is an object in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ with structures arising from the ones of $A \otimes H$, ie.,

$$\begin{aligned} & (a \otimes h \otimes m)b = a\beta^{-1}(b_{[0]}) \otimes h\alpha^{-1}(b_{[1]}) \otimes \mu(m), \\ & \rho_{A\otimes H\otimes M}(a \otimes h \otimes m) = \beta^{-1}(a) \otimes h_{(1)} \otimes \mu^{-1}(m) \otimes \alpha^2(h_{(2)}), \end{aligned}$$

for all $h \in H, a, b \in A$ and $m \in M$. It is routine to check that $A \otimes H \otimes M$ is a right (H, α)-Hom-comodule and a right (A, β)-Hom-module. Now we only check the compatibility condition. Indeed, for any $a, b \in A$, we have

$$\begin{aligned}
& \rho_{A\otimes H\otimes M}((a \otimes h \otimes m)b) \\
= \ & \rho_{A\otimes H\otimes M}(a\beta^{-1}(b_{[0]}) \otimes h\alpha^{-1}(b_{[1]}) \otimes \mu(m)) \\
= \ & \beta^{-1}(a)\beta^{-2}(b_{[0]}) \otimes h_{(1)}\alpha^{-1}(b_{[1](1)}) \otimes m \otimes \alpha^2(h_{(2)})\alpha(b_{[1](2)}) \\
= \ & \beta^{-1}(a)\beta^{-1}(b_{[0][0]}) \otimes h_{(1)}\alpha^{-1}(b_{[0][1]}) \otimes m \otimes \alpha^2(h_{(2)})b_{[1]} \\
= \ & [\beta^{-1}(a) \otimes h_{(1)} \otimes \mu^{-1}(m)]b_{[0]} \otimes \alpha^2(h_{(2)})b_{[1]} \\
= \ & \rho_{A\otimes H\otimes M}(a \otimes h \otimes m)b,
\end{aligned}$$

as desired. Next we will prove that f is a k-split surjection. Let $g : M \to A\otimes H\otimes M, g(m) = 1_A \otimes \alpha^{-1}(m_{[1]}) \otimes m_{[0]}$, for all $m \in M$. Then g is right (H, α)-colinear and for any $m \in M$, we have

$$\begin{aligned}
(f \circ g)(m) &= f(1_A \otimes \alpha^{-1}(m_{[1]}) \otimes m_{[0]}) \\
&= \mu(m_{[0][0]})\gamma(m_{[0][1]})(\alpha^{-1}(m_{[1]})) \\
&= m_{[0]}\gamma(m_{[1](1)})(m_{[1](2)}) \\
&= m.
\end{aligned}$$

For all $a, b \in A$ and $h \in H, m \in M$, we have

$$\begin{aligned}
& f((a \otimes h \otimes m)b) \\
= \ & f(a\beta^{-1}(b_{[0]}) \otimes h\alpha^{-1}(b_{[1]}) \otimes \mu(m)) \\
= \ & \mu^2(m_{[0]})\gamma(m_{[1]})(\alpha^{-2}(h\alpha^{-1}(b_{[1]}))S^{-1}(\alpha^{-1}(a_{[1]}\alpha^{-1}(b_{[0][1]}))))\beta(a_{[0]}\beta^{-1}(b_{[0][0]})) \\
= \ & \mu^2(m_{[0]})\gamma(m_{[1]})(\alpha^{-2}(h)\alpha^{-3}(b_{[1]})S^{-1}(\alpha^{-2}(b_{[0][1]}))S^{-1}(\alpha^{-1}(a_{[1]})))[\beta(a_{[0]})b_{[0][0]}] \\
= \ & \mu^2(m_{[0]})\gamma(m_{[1]})(\alpha^{-1}(h)\{[\alpha^{-3}(b_{[1](2)})S^{-1}(\alpha^{-3}(b_{[1](1)}))]S^{-1}(\alpha^{-1}(a_{[1]}))\})[\beta(a_{[0]})\beta^{-1}(b_{[0]})] \\
= \ & \mu^2(m_{[0]})\gamma(m_{[1]})(\alpha^{-1}(h)S^{-1}(a_{[1]}))[\beta(a_{[0]})\beta^{-2}(b)] \\
= \ & \mu^2(m_{[0]})\{[\gamma(\alpha^{-1}(m_{[1]}))(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))\beta(a_{[0]})]\beta^{-1}(b)\} \\
= \ & \{\mu(m_{[0]})[\gamma(\alpha^{-1}(m_{[1]}))(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))\beta(a_{[0]})]\}b \\
= \ & f(a \otimes h \otimes m)b.
\end{aligned}$$

Then f is a right (A, β)-linear. It remains to prove that f is also right (H, α)-colinear. For this purpose, we do the following calculations:

$$\begin{aligned}
& (f \otimes id_H)\rho_{A\otimes H\otimes M}(a \otimes h \otimes m) \\
= \ & (f \otimes id_H)(\beta^{-1}(a) \otimes h_{(1)} \otimes \mu^{-1}(m) \otimes \alpha^2(h_{(2)})) \\
= \ & f(\beta^{-1}(a) \otimes h_{(1)} \otimes \mu^{-1}(m)) \otimes \alpha^2(h_{(2)})) \\
= \ & m_{[0]}\gamma(\alpha^{-2}(m_{[1]}))(\alpha^{-2}(h_{(1)})S^{-1}(\alpha^{-2}(a_{[1]})))a_{[0]} \otimes \alpha^2(h_{(2)})).
\end{aligned}$$

On the other side,

$$\begin{aligned}
& \rho_{A\otimes H\otimes M}\circ f(a\otimes h\otimes m)\\
= \;& \rho_{A\otimes H\otimes M}(\mu(m_{[0]})\gamma(\alpha^{-1}(m_{[1]}))(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))\beta(a_{[0]}))\\
= \;& \mu(m_{[0][0]})\gamma(\alpha^{-1}(m_{[1]}))(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))_{[0]}\beta(a_{[0][0]})\\
& \otimes\mu(m_{[0][1]})\gamma(\alpha^{-1}(m_{[1]}))(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))_{[1]}\beta(a_{[0][1]})\\
= \;& m_{[0]}\gamma(m_{[1](2)})(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))_{[0]}\beta(a_{[0][0]})\\
& \otimes\mu(m_{[1](1)})\gamma(m_{[1](2)})(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))_{[1]}\beta(a_{[0][1]})\\
= \;& m_{[0]}\gamma(m_{[1](2)})(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))_{[0]}\beta(a_{[0][0]})\\
& \otimes[m_{[1](1)}\gamma(m_{[1](2)})(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))_{[1]}]\beta^2(a_{[0][1]})\\
\overset{(4.1)}{=} \;& m_{[0]}\gamma(\alpha^{-2}(m_{[1]}))(\alpha^{-2}(h_{(1)})S^{-1}(\alpha^{-1}(a_{[1](2)})))\beta(a_{[0][0]})\\
& \otimes\alpha(\alpha^{-1}(h_{(2)})S^{-1}(a_{[1](1)}))\alpha^2(a_{[0][1]})\\
= \;& m_{[0]}\gamma(\alpha^{-2}(m_{[1]}))(\alpha^{-2}(h_{(1)})S^{-1}(a_{[1](2)(2)}))a_{[0]})\\
& \otimes(h_{(2)}S^{-1}(\alpha^2(a_{[1](2)(1)})))\alpha^2(a_{[1](1)})\\
= \;& m_{[0]}\gamma(\alpha^{-2}(m_{[1]}))(\alpha^{-2}(h_{(1)})S^{-1}(\alpha^{-1}(a_{[1](2)})))a_{[0]})\\
& \otimes\alpha(h_{(2)})(S^{-1}(\alpha^2(a_{[1](1)(2)}))\alpha^2(a_{[1](1)(1)}))\\
= \;& m_{[0]}\gamma(\alpha^{-2}(m_{[1]}))(\alpha^{-2}(h_{(1)})S^{-1}(\alpha^{-2}(a_{[1]})))a_{[0]})\otimes\alpha^2(h_{(2)}).
\end{aligned}$$

Hence f is an epimorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ and has a (H,α)-colinear section. For any k-free presentation of (M,μ) in the category of k-modules $k^I\overset{\pi}{\to}M\to 0$, there is an epimorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$:

$$(A|oH)^I\cong A\otimes H\otimes k\overset{g}{\to}M\to 0,$$

where $g=f\circ(id_A\otimes id_H\otimes\pi)$. Hence $A\otimes H$ is a generator in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. The proof is completed. □

5 The affineness criterion for relative Hom-Hopf modules

Proposition 5.1. Let (H,α) be a monoidal Hom-Hopf algebra with a bijective antipode and (A,β) a right (H,α)-Hom-comodule algebra. Assume that there exists a total quantum integral $\gamma:H\to Hom(H,A)$. Then $\widetilde{\rho}:A\to A\otimes H,\widetilde{\rho}(a)=a_{[0]}\otimes a_{[1]}$ splits in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$.

Proof. We consider the map

$$\lambda:A\otimes H\to A\qquad\qquad\lambda(a\otimes h)=a_{[0]}\gamma(a_{[1]})(h)$$

for all $a\in A$ and $h\in H$ is a right (H,α)-colinear retraction of $\widetilde{\rho}$. In particular, by Eqs.(5.1) and 5.2, we obtain $\lambda(1_A\otimes 1_H)=1_A$, and

$$\lambda(\beta^{-1}(a)\otimes h_{(1)})\otimes h_{(2)}=\lambda(a\otimes h)_{[0]}\otimes\lambda(a\otimes h)_{[1]}. \tag{5. 1}$$

We define now

$$\Lambda : A \otimes H \to A \qquad \Lambda(a \otimes h) = \lambda(1_A \otimes \alpha^{-1}(h)S^{-1}(a_{[1]}))\beta(a_{[0]})$$

for all $h \in H$ and $a \in A$. Then, for all $a \in A$ we have

$$\begin{aligned}(\Lambda \circ \widetilde{\rho})(a) &= \Lambda(a_{[0]} \otimes a_{[1]}) \\ &= \lambda(1_A \otimes \alpha^{-1}(a_{[1]})S^{-1}(a_{[0][1]}))\beta(a_{[0][0]}) \\ &= \lambda(1_A \otimes a_{[1](2)}S^{-1}(a_{[1](1)}))a_{[0]} \\ &= \lambda(1_A \otimes 1_H)\beta^{-1}(a) = a,\end{aligned}$$

i.e., Λ is still a retraction of $\widetilde{\rho}$. Now, for all $h \in H, a, b \in A$, we have

$$\begin{aligned}\Lambda((a \otimes h)b) &= \Lambda(ab_{[0]} \otimes hb_{[1]}) \\ &= \lambda(1_A \otimes \alpha^{-1}(hb_{[1]})S^{-1}(a_{[1]}b_{[0][1]}))\beta(a_{[0]}b_{[0][0]}) \\ &= \lambda(1_A \otimes h[\alpha^{-1}(b_{[1](2)})S^{-1}(\alpha^{-1}(b_{[1](1)}))]S^{-1}(a_{[1]}))\beta(a_{[0]})b_{[0][0]} \\ &= \lambda(1_A \otimes hS^{-1}(\alpha(a_{[1]})))[\beta(a_{[0]})\beta^{-1}(b)] \\ &= \lambda(1_A \otimes \alpha^{-1}(h)S^{-1}(a_{[1]}))\beta(a_{[0]})b \\ &= \Lambda(a \otimes h)b\end{aligned}$$

Hence, Λ is right (A, β)-linear. It remains to prove that Λ is also (H, α)-colinear, we have

$$\begin{aligned}\widetilde{\rho}\Lambda(a \otimes h) &= \widetilde{\rho}(\lambda(1_A \otimes \alpha^{-1}(h)S^{-1}(a_{[1]}))\beta(a_{[0]})) \\ &= \lambda(1_A \otimes \alpha^{-1}(h)S^{-1}(a_{[1]}))_{[0]}\beta(a_{[0][0]}) \otimes \lambda(1_A \otimes \alpha^{-1}(h)S^{-1}(a_{[1]}))_{[1]}\alpha(a_{[0][1]}) \\ &\overset{(5.1)}{=} \lambda(1_A \otimes \alpha^{-1}(h_{(1)})S^{-1}(a_{[1](2)}))\beta(a_{[0][0]}) \otimes [\alpha^{-1}(h_{(2)})S^{-1}(a_{[1](1)})]\alpha(a_{[0][1]}) \\ &= \lambda(1_A \otimes \alpha^{-1}(h_{(1)})S^{-1}(a_{[1](2)}))\beta(a_{[0][0]}) \otimes h_{(2)}[S^{-1}(a_{[1](1)})a_{[0][1]}] \\ &= \lambda(1_A \otimes \alpha^{-1}(h_{(1)})S^{-1}(\alpha(a_{[1](2)(2)})))a_{[0]} \otimes h_{(2)}[S^{-1}(\alpha(a_{[1](2)(1)}))a_{[1](1)}] \\ &= \lambda(1_A \otimes \alpha^{-1}(h_{(1)})S^{-1}(a_{[1](2)}))a_{[0]} \otimes h_{(2)}[S^{-1}(\alpha(a_{[1](1)(2)}))\alpha(a_{[1](1)(1)})] \\ &= \lambda(1_A \otimes \alpha^{-1}(h_{(1)})S^{-1}(\alpha^{-1}(a_{[1]})))a_{[0]} \otimes \alpha(h_{(2)}) \\ &= (\Lambda \otimes id_H)\rho_{A \otimes H},\end{aligned}$$

i.e., proved that Λ is a retraction of $\widetilde{\rho}$ in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. □

Recall from [12] that the coinvariants of A as

$$B = A^{coH} = \{a \in A | \widetilde{\rho}(a) = \beta^{-1}(a) \otimes 1_H\}.$$

Then B is a subalgebra of A.

Proposition 5.2. Let (H, α) be a monoidal Hom-Hopf algebra with a bijective antipode and (A, β) a right (H, α)-Hom-comodule algebra. Assume that there exists $\gamma : H \to Hom(H, A)$ a total quantum integral. Then

(1) B is a direct summand of A as a left B-submodule,

(1) B is a direct summand of A as a right B-submodule.

Proof. We shall prove that there exists a well defined left trace given by the formula

$$t^l : (A, \beta) \to (B, \beta) \qquad t^l(a) = \lambda(a \otimes 1_H) = a_{[0]}\gamma(a_{[1]})(1_H)$$

for all $a \in A$, by Eq.(5.1) we obtain $\widetilde{\rho}(t^l(a)) = t^l(a) \otimes 1_H$, ie., $t^l(a) \in B$, for all $a \in A$. Now for all $b \in B$ and $a \in A$, we have

$$\begin{aligned} t^l(ba) &= b_{[0]}a_{[0]})\gamma(b_{[1]}a_{[1]})(1_H) \\ &= [ba_{[0]}]\gamma(a_{[1]})(1_H) \\ &= ba_{[0]}\gamma(a_{[1]})(1_H) \\ &= bt^l(a), \end{aligned}$$

hence t^l is a left (B, β)-module map and finally

$$t^l(1_A) = 1_A\gamma(1_H)(1_H) = 1_A.$$

hence t^l is a left (B, β)-module retraction of the inclusion $B \subset A$.

(2) Similarly, we can prove that the map given by the formula

$$t^r : (A, \beta) \to (B, \beta), \qquad t^r(a) = \Lambda(a \otimes 1_H) = \gamma(1_H)(S^{-1}(\alpha^2(a_{[1]})))\beta(a_{[0]})$$

for all $a \in A$, t^r is a right (B, β)-module retraction of the inclusion $B \subset A$. □

Definition 5.3. Let (H, α) be a monoidal Hom-Hopf algebra with a bijective antipode and (A, β) a right (H, α)-Hom-comodule algebra. Assume that there exists $\gamma : H \to Hom(H, A)$ a total quantum integral. The map

$$t^l : (A, \beta) \to (B, \beta), \qquad t^l(a) = a_{[0]}\gamma(a_{[1]})(1_H)$$

for all $a \in A$ is called the quantum trace associated of γ.

Now, we shall construct functors connecting $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ and $\widetilde{\mathscr{H}}(\mathscr{M}_k)_B$. First, if $(M, \mu) \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$, then

$$M^{coH} = \{m \in M | \widetilde{\rho}(m) = \mu^{-1}(m) \otimes 1_H\}.$$

is the right B-module of the coinvariants of M. Furthermore, $M \to M^{coH}$ gives us a covariant functor

$$(-)^{coH} : \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H \to \widetilde{\mathscr{H}}(\mathscr{M}_k)_B.$$

Now, for $(N, \nu) \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_B$, $N \otimes_B A \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ via the structures

$$\begin{aligned} (a \otimes_B n)a' &= a\beta^{-1}(a') \otimes_B \nu(n) \\ \rho_{A\otimes_B N}(a \otimes_B n) &= a_{[0]} \otimes \nu^{-1}(n) \otimes \alpha(a_{[1]}) \end{aligned}$$

for all $n \in N, a, a' \in A$. In this way, we have constructed a covariant functor called the induction functor

$$A \otimes_B - : \widetilde{\mathscr{H}}(\mathscr{M}_k)_B \to \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H.$$

Proposition 5.4. Let (H, α) be a monoidal Hom-Hopf algebra with a bijective antipode and (A, β) a right (H, α)-Hom-comodule algebra. Then the induction functor $A \otimes_B - : \widetilde{\mathscr{H}}(\mathscr{M}_k)_B \to \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ is a left adjoint of the coinvariant functor $(-)^{coH} : \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H \to \widetilde{\mathscr{H}}(\mathscr{M}_k)_B$.

Proof. Similar to [17]. □

Recall from [12] that $A \otimes H \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ with the structures

$$\begin{cases} (a \otimes h) \cdot b = ab_{[0]} \otimes hb_{[1]}; \\ \rho_{A\otimes H}(a \otimes h) = (\beta^{-1}(a) \otimes h_{(1)}) \otimes \alpha(h_{(2)}), \end{cases}$$

for any $a, b \in A$ and $h \in H$, and we identify $(A \otimes H)^{coH} \cong A$ via $a \otimes 1_H \mapsto a$. Then the the adjunction map can be viewed as a map in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$, as

$$\psi : A \otimes_B A \to A \otimes H, \quad \psi(a \otimes b) = ab_{[0]} \otimes b_{[1]}$$

for all $a, b \in A$. Here $A \otimes_B A \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ with the structures

$$\begin{cases} (a \otimes_B b) \cdot a' = \beta(a) \otimes_B b\beta^{-1}(a'); \\ \rho_{A\otimes_B A}(a \otimes_B b) = (\beta^{-1}(a) \otimes b_{[0]}) \otimes \alpha(b_{[1]}), \end{cases}$$

for all $a, a', b \in A$.

Definition 5.5. Let (H, α) be a monoidal Hom-Hopf algebra with a bijective antipode and (A, β) a right (H, α)-Hom-comodule algebra, and $B = A^{coH}$. Then A/B is called a Galois extension if the the canonical map

$$\psi : A \otimes_B A \to A \otimes H, \quad \psi(a \otimes b) = \beta^{-1}(a)b_{[0]} \otimes \alpha(b_{[1]})$$

is bijective.

Theorem 5.6. Let (H, α) be a monoidal Hom-Hopf algebra with a bijective antipode and (A, β) a right (H, α)-Hom-comodule algebra, and $B = A^{coH}$. Assume that there exists $\gamma : H \to Hom(H, A)$ a total quantum integral. Then

$$\eta_N : N \to (A \otimes_B N)^{coH}, \qquad \eta_N(n) = 1_A \otimes_B n$$

is an isomorphism of right (B, β)-modules for all $(N, \nu) \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_B$.

Proof. Using the left quantum $t^l : (A, \beta) \to (B, \beta)$ we shall construct an inverse of η_N, we define

$$\theta_N : (A \otimes_B N)^{coH} \to N, \qquad \theta_N(a_i \otimes_B n_i) = \sum t^l(a_i)n_i$$

for any $a_i \otimes_B n_i \in (A \otimes_B N)^{coH}$. Since $t^l(1_A) = 1_A$, and it is easy to check that $\theta_N \circ \eta_N = id_N$. Now, let $a_i \otimes_B n_i \in (A \otimes_B N)^{coH}$. Then

$$\alpha^{-1}(a_i) \otimes_B \nu^{-1}(n_i) \otimes 1_H = a_{[0]} \otimes_B \nu^{-1}(n_i) \otimes \alpha(a_{[1]}).$$

It follows that

$$\alpha^{-1}(a_i) \otimes_B \nu^{-1}(n_i) \otimes 1_A = a_{[0]} \otimes_B \nu^{-1}(n_i) \otimes \gamma(a_{[1]})(1_H).$$

Furthermore, we have

$$\alpha^{-1}(a_i)1_A \otimes_B \nu^{-1}(n_i) = a_{[0]}\gamma(a_{[1]})(1_H) \otimes_B \nu^{-1}(n_i).$$

then, we have

$$a_i \otimes_B n_i = t^l(a_i) \otimes_B n_i.$$

Hence, we have

$$(\eta_N \circ \theta_N)(a_i \otimes_B n_i) = \sum 1_A \otimes_B t^l(a_i) n_i = \sum t^l(a_i) \otimes_B n_i = a_i \otimes_B n_i,$$

Then θ_N is an inverse of η_N. □

We will now prove the main result of this section, that is, the affineness criterion for relative Hom-Hopf modules.

Theorem 5.7. Let (H, α) be a monoidal Hom-Hopf algebra with a bijective antipode and (A, β) a right (H, α)-Hom-comodule algebra, and $B = A^{coH}$. Assume that

(1) there exists a total quantum integral $\gamma : H \to Hom(H, A)$;

(2) the canonical map

$$\beta : A \otimes_B A \to A \otimes H, \quad a \otimes_B b \mapsto \beta^{-1}(a) b_{[0]} \otimes \alpha(b_{[1]})$$

is surjective. Then the induction functor $A \otimes_B - : \widetilde{\mathscr{H}}(\mathscr{M}_k)_B \to \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ is an equivalence of categories.

Proof. In Theorem 5.6 we have shown that the adjunction map $\eta_N : N \to (A \otimes_B N)^{coH}$ is an isomorphism for all $(N, \nu) \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_B$. It remains to prove that the other adjunction map, namely $\beta_M : M^{coH} \otimes_B A \to M$ is also an isomorphism for all $(M, \mu) \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$.

Let (V, ω) be a k-module. Then $A \otimes V \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ via the structures induced by A, i.e.,

$$(a \otimes v)a' = a\beta^{-1}(a') \otimes \omega(v)$$
$$\rho_{A \otimes V}(a \otimes v) = a_{[0]} \otimes \omega^{-1}(v) \otimes \alpha(a_{[1]})$$

for all $a, b \in A$ and $v \in V$. In particular, for $V = A, A \otimes A \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ via

$$(a \otimes b) \cdot a' = a\beta^{-1}(a') \otimes \beta(b); \tag{5. 2}$$
$$\rho_{A \otimes A}(a \otimes b) = a_{[0]} \otimes \beta^{-1}(b) \otimes \alpha(a_{[1]}) \tag{5. 3}$$

for all $a, b, a' \in A$. We will first prove that the adjunction map $\beta_{A \otimes V} : (A \otimes V)^{coH} \otimes_B A \to A \otimes V$ is an isomorphism for any k-module V.

First, $V \otimes B$ and $B \otimes V \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_B$ via the usual B-actions $(v \otimes a) \cdot b = \omega(v) \otimes a\beta^{-1}(b)$, and $a' \cdot (b' \otimes v') = \beta^{-1}(a')b' \otimes \omega(v')$ for all $a, b, a', b' \in B$ and $v, v' \in V$. The flip map $\tau : V \otimes B \to B \otimes V, \quad \tau(v \otimes b) = b \otimes v$, for all $b \in B$ and $v \in V$, is an isomorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_B$. On the other hand $A \otimes V \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ via the structures induced by A, i.e.

$$(v \otimes a) \cdot b = \omega(v) \otimes a\beta^{-1}(b),$$
$$\rho_{V \otimes A}(v \otimes a) = \omega^{-1}(v) \otimes a_{[0]} \otimes \alpha(a_{[1]}),$$

It is easy to see that the flip map $\tau : A \otimes V \to V \otimes A, \quad \tau(a \otimes v) = v \otimes a$ is an isomorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$.

Applying Theorem 5.6 for $N = V \otimes B \cong B \otimes V$, we obtain the following isomorphisms in $\mathcal{M}_B$:

$$B \otimes V \cong V \otimes B \cong (V \otimes B \otimes_B A)^{coH} \cong (V \otimes A)^{coH} \cong (A \otimes V)^{coH}.$$

Hence, $(A \otimes V)^{coH} \otimes_B A \cong A \otimes V$.

Let us define

$$\widetilde{\psi} : A \otimes_B A \to A \otimes H, \quad a \otimes_B b \mapsto \beta^{-1}(a)b_{[0]} \otimes \alpha(b_{[1]}),$$

for all $a, b \in A$. As ψ is surjective, $\widetilde{\psi}$ is surjective, because $\widetilde{\psi} = \psi \circ can$, where $can : A \otimes A \to A \otimes_B A$ is the canonical surjection.

Let us define now

$$\xi : A \otimes A \to A \otimes H, \quad \xi(a \otimes b) = (\widetilde{\psi} \circ \tau)(a \otimes b) = \beta^{-1}(b)a_{[0]} \otimes \alpha(a_{[1]}),$$

for all $a, b \in A$. The map ξ is surjective, as $\widetilde{\psi}$ and τ are. We will prove that ξ is a morphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. where $A \otimes A$ and $A \otimes H$ are realtive Hom-Hopf modules, respectively. Indeed,

$$\begin{aligned} \xi((a \otimes b)c) = \xi(a\beta^{-1}(c) \otimes \beta(b)) &= b[a_{(-1)}\beta^{-1}(c_{(-1)})] \otimes \alpha(a_{(1)})(c_{(1)}) \\ &= [\beta^{-1}(b)a_{(-1)}]c_{(-1)} \otimes \alpha(a_{(1)})(c_{(1)}) \\ &= \xi(a \otimes b)c, \end{aligned}$$

and

$$\begin{aligned} \rho_{A\otimes H}\xi(a \otimes b) &= \rho_{A\otimes H}(\beta^{-1}(b)a_{[0]} \otimes \alpha(a_{[1]})) \\ &= [\beta^{-2}(b)\beta^{-1}(a_{[0]}) \otimes \alpha(a_{[1](1)})] \otimes \alpha^2(a_{[1](2)}) \\ &= [\beta^{-2}(b)a_{[0]} \otimes \alpha(a_{[0][1]})] \otimes \alpha(a_{[1]}) \\ &= (\xi \otimes id)(a_{[0]} \otimes \beta^{-1}(b) \otimes \beta^{-1}(a_{[1]})) \\ &= (\xi \otimes id_H)\rho_{A\otimes A}(a \otimes b). \end{aligned}$$

Hence, ξ is a surjective morphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. $A \otimes H$ is projective as a left A-module, where $A \otimes H$ is a left A-module in the usual way. On the other hand, by Proposition 3.1 we obtain the map

$$u : A \otimes H \to A \otimes H, \quad a \otimes h \mapsto \beta(a_{[0]}) \otimes h\alpha^{-1}(a_{[1]})$$

is an isomorphism of left A-modules. It follows that there exits $\zeta : A \otimes H \to A \otimes A$ such that $\xi \circ \zeta = id_{H\boxtimes A}$ since $A \otimes A \to A \otimes H$ is surjective. Hence, ξ splits in the category of right A-modules. In particular ξ is a k-split epimorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$.

Let now $(M, \mu) \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. Then $A \otimes A \otimes M \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ via the structures arising from $A \otimes A$, that is,

$$(a \otimes b \otimes m) \cdot c = a\beta^{-2}(c) \otimes \beta(b) \otimes \mu(m); \tag{5. 4}$$

$$\rho_{A\otimes A\otimes M}(a \otimes b \otimes m) = \beta^{-1}(a) \otimes b_{[0]} \otimes \mu^{-1}(m) \otimes \alpha^2(b_{[1]}), \tag{5. 5}$$

for all $a,b,c \in A$ and $m \in M$. On the other hand, $A \otimes H \otimes M \in \widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ via the structures arising from $A \otimes H$, that is,

$$(a \otimes h \otimes m)b = a\beta^{-1}(b_{[0]}) \otimes h\alpha^{-1}(b_{[1]}) \otimes \mu(m)$$
$$\rho_{A\otimes H\otimes M}(a \otimes h \otimes m) = \beta^{-1}(a) \otimes h_{(1)} \otimes \mu^{-1}(m) \otimes \alpha^2(h_{(2)}).$$

for all $a,b \in A, h \in H$ and $m \in M$. We obtain that

$$\xi \otimes id_M : A \otimes A \otimes M \to A \otimes H \otimes M$$

is a k-split epimorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$.

By Theorem 4.8 we obtain that the map

$$f : A \otimes H \otimes M \to M, \quad a \otimes h \otimes m \mapsto \mu(m_{[0]})\gamma(\alpha^{-1}(m_{[1]}))(\alpha^{-2}(h)S^{-1}(\alpha^{-1}(a_{[1]})))\beta(a_{[0]})$$

is a k-split epimorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. Hence, the composition

$$g = f \circ (\xi \otimes id_M) : A \otimes A \otimes M \to M,$$
$$a \otimes b \otimes m \mapsto \mu(m_{[0]})\gamma(\alpha^{-1}(m_{[1]}))(S^{-1}(b_{[1]}))b_{[0]}\beta^{-1}(a)$$

is a k-split epimorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$. We note that the structure of $A \otimes A \otimes M$ as an object in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$ is of the form $A \otimes V$, for the k-module $V = A \otimes M$.

To conclude, we have constructed a k-split epimorphism in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$

$$A \otimes A \otimes M = (M_1, \pi) \xrightarrow{g} (M, \mu) \longrightarrow 0$$

such that the adjunction map ψ_{M_1} for (M_1, π) is bijective. As g is k-split and there exists a total quantum integral $\gamma : H \to Hom(H, A)$, we obtain that g also splits in $\widetilde{\mathscr{H}}(\mathscr{M}_k)^H$. In particular, the sequence

$$(M_1^{coH}, \pi) \xrightarrow{g^{coH}} (M^{coH}, \mu) \longrightarrow 0$$

is exact. Continuing the resolution with $Ker(g)$ instead of M, we obtain an exact sequence in $\widetilde{\mathscr{H}}(\mathscr{M}_k)_A^H$

$$(M_2, P) \longrightarrow (M_1, \pi) \longrightarrow (M, \mu) \longrightarrow 0$$

which splits in $\widetilde{\mathscr{H}}(\mathscr{M}_k)^H$ and the adjunction maps for (M_1, π) and (M_2, P) are bijective. Using the Five lemma we obtain that the adjunction map for (M, μ) is bijective. □

Finally we consider a special case. In this case if setting $A = H$, then (A, β) is a right H-comodule algebra in a natural way, and we can define the coinvariants of (H, α) as

$$B = H^{coH} = \{h \in H | \tilde{\rho}(h) = \alpha^{-1}(h) \otimes 1_H\}$$

then (B, α) is a subalgebra of (H, α). Hence we can obtain the following result.

Corollary 5.8. Let (H, α) be a monoidal Hom-Hopf algebra with a bijective antipode and (A, β) a right (H, α)-Hom-comodule algebra, and $B = H^{coH}$. Assume that
(1) there exists a total quantum integral $\gamma : H \to Hom(H, H)$;
(2) the canonical map

$$\psi : H \otimes_B H \to H \otimes H, \quad h \otimes_B g \mapsto \alpha^{-1}(h)g_{(1)} \otimes \alpha(g_{(2)})$$

is surjective. Then the induction functor $- \otimes_B H : \widetilde{\mathscr{H}}(\mathscr{M}_k)_B \to \widetilde{\mathscr{H}}(\mathscr{M}_k)_H^H$ is an equivalence of categories.

Acknowledgements

The work is supported by the NSF of Jiangsu Province (No. BK2012736) and the Fund of Science and Technology Department of Guizhou Province (No. 2014GZ81365).

References

[1] S. Caenepeel and I. Goyvaerts, Monoidal Hom-Hopf algebras. *Comm. Algebra* **39** (2011) 2216-2240.

[2] Y. Y. Chen, Z. W. Wang and L. Y. Zhang, Integrals for monoidal Hom-Hopf algebras and their applications. *J. Math. Phy.* **54**(7) (2013) 073515.

[3] Y. Y. Chen, Z. W. Wang and L. Y. Zhang, The FRT-type theorem for the Hom-Long equation. *Comm. Algebra* **41** (2013) 3931-3948.

[4] Y. Y. Chen and L. Y. Zhang. The category of Yetter-Drinfel'd Hom-modules and the quantum Hom-Yang-Baxter equation. *J. Math. Phys.* **55**(3) (2014) 031702.

[5] E. Cline, B. Parshall, L. Scott, Induced modules and affine quotient. *Math. Ann.* **230** (1977) 1-14.

[6] Y. Doi, On the structure of relative Hopf modules. *Comm. Algebra* **11**(3) (1983) 243-255.

[7] Y. Doi, Algebras with total integrals. *Comm. Algebra* **13** (1985) 2137-2159.

[8] Y. Doi, Hopf extensions of algebras and Maschke type theorems. *Israel J. Math.* **72** (1990) 99-108.

[9] Y. Frégier and A. Gohr, On Hom type algebras. *J. Gen. Lie Theory Appl.* **4**(2010) 16.

[10] Y. Frégier, A. Gohr and S. D. Silvestrov, Unital algebras of Hom-associative type and surjective or injective twistings. *J. Gen. Lie Theory Appl.* **3** (2009) 285-295.

[11] A. Gohr, On Hom-algebras with surjective twisting. *J. Algebra* **324** (2010) 1483-1491.

[12] S. J. Guo and X. L. Chen, A Maschke type theorem for relative Hom-Hopf modules. Accepted by *Czech. Math. J.* 2014.

[13] J. T. Hartwig, D. Larsson and S. D. Silvestrov, Deformations of Lie algebras using σ-derivations. *J. Algebra* **295** (2006) 314-361.

[14] A. Makhlouf and S. D. Silvestrov, Hom-algebra stuctures. *J. Gen. Lie Theory Appl.* **2** (2008) 51-64.

[15] A. Makhlouf, S. Silvestrov. Hom-Lie admissible Hom-coalgebras and Hom-Hopf algebras. *J. Gen. Lie Theory in Mathematics, Physics and beyond*. Springer-Verlag, Berlin, 2009, pp. 189-206.

[16] A. Makhlouf and S. D. Silvestrov, Hom-algebras and Hom-coalgebras. *J. Algebra Appl.* **9** (2010) 553-589.

[17] C. Menini and G. Militaru, Integral, quantum Galois extensions, and the affineness criterion for quantum Yetter-Drinfeld modules. *J. Algebra* **247** (2002) 467-508.

[18] U. Oberst, Affine Quotientenschemata nach affinen, algebraischen Gruppen und induzierte Darstellungen. *J. Algebra* **44**(1977) 503-538.

[19] S. X. Wang and S. J. Guo. Symmetries and the u-condition in Hom-Yetter-Drinfeld categories. *J. Math. Phys.* **55** (2014) 081708.

[20] D. Yau, Hom-algebras and homology. *J. Lie Theory* **10** (2009) 409-421.

[21] D. Yau, Hom-bialgebras and comodule algebras. *Int. Electron. J. Algebra* **8** (2010) 45-64.

[22] D. Yau. The Hom-Yang-Baxter equation, Hom-Lie algebras, and quasi-triangular bialgebras. *J. Phys.* **42**(16)(2009) 165202.